

Powered by

LINDNER
Feed & Milling

www.lindnerfeed.com

*Feeding
Guide*

Powered by

LINDNER

Lindner Feed & Milling is a fourth generation ranch and farm feed and supply company located in the central Texas town of Comfort.

We provide high-quality feeds and ranching products at low prices to farmers, ranchers, and stock show participants in surrounding communities.

Lindner Feed was founded in 1938 by Michael Lindner. His son, Percy Linder, operated the mill in the 1950s and 1960s. Today, Lindner Feed is managed by his grandchildren and great-grandchildren. The Mill, as it is affectionately known, is a fixture in the Comfort, Texas landscape and business community. We are proud supporters of many youth programs including 4-H, FFA, and the Kendall and Kerr County Junior Livestock Shows.

We believe that quality does matter. Lindner Feed is committed to our customers and focused on their results.



Lindner Feed is proud to announce the recent addition of a fully computerized state-of-the-art pellet mill to our facilities. With this new machinery, we are now capable of producing commercial and custom pelleted feed in a variety of sizes.

Our stock show pig feeds are known across the state. We are proud to congratulate the many exhibitors who have found success using our feeds. If you have a winner that was powered by Lindner, we'd love to hear about it.

Our motto is "Committed to our Customers...Focused on Results!" If we can assist you in any way, please feel free to contact us at any time. Inquiries and comments can be directed to Conda Maze at 281-642-4017.

LINDNER 610 EARLY WEAN 22 MX BN MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibriotic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox.....0.0055%(50 grams per ton)
Pyrantel Tartrate.....0.0106% (96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein-Min.....22.0%	Lysine-Min.....1.73%
Crude Fat-Min.....5.6%	Crude Fiber-Max.....2.4%
Calcium-Min.....0.8%	Calcium-Max.....1.0%
Phosphorous-Min.....0.64%	Salt-Min.....None added
Selenium-Min.....0.4 ppm	Zinc-Min.....3700 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrus Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Early wean 22% is a complete feed specifically formulated for 8 to 15lb. pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 15lb. body weight. Then they should be fed an approved starter ration.

WARNING: Do not feed to swine weighing more than 75lb. body weight. Do not feed to swine within 10 weeks of slaughter

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

Key Information

Protein: 22%

Fat: 6%

Lysine: 1.75%

Medication: MXBN

Withdrawal: 10 weeks

Feed to pigs weighing 8lbs. to 14lbs.

LINDNER 611 Scud Missile- PIG STARTER 23% MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For increased rate of gain and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... 50 grams per ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	23.0%	Lysine-Min.....	1.42%
Crude Fat-Min.....	7.10%	Crude Fiber-Max.....	4.20%
Calcium-Min.....	0.78%	Calcium-Max.....	0.85%
Phosphorous-Min.....	0.67%	Salt-Min.....	0.01%
Selenium-Min.....	0.3 ppm	Zinc-Min.....	3200 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Soybean Hulls, Dried Whey, Yeast Culture, Plasma, Animal fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

611 Scud Missile is a complete feed specifically formulated for 8 to 15 lb. pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 15lb. body weight. Then they should be fed an approved starter ration.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

Key Information

Protein: 23%

Fat: 7.10%

Lysine: 1.42%

Medication: Chlortetracycline

Withdrawal: none

Feed to pigs weighing 8lbs. to 15lbs.

LINDNER 612 SCUD MISSILE - PIG STARTER 23% MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For the controle of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella chloeraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Tiamulin hydrogen fumarate 35 g/ton Chlortetracycline.....400 grams per ton

GUARANTEED ANALYSIS

Crude Protein-Min	23.0%	Lysine-Min.....	1.42%
Crude Fat-Min.....	7.10%	Crude Fiber-Max.....	4.20%
Calcium-Min.....	0.78%	Calcium-Max.....	0.85%
Phosphorous-Min.....	0.67%	Salt-Min.....	0.01%
Selenium-Min.....	0.3 ppm	Zinc-Min.....	3200 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Soybean Hulls, Dried Whey, Yeast Culture, Plasma, Animal fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysine, Copper Sulfate, Artificial flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

612 Scud Missile is a complete feed specifically formulated for 8 to 15 lb. pigs fed for exhibition. This product must be fed continuously as the sole ration for 14 days only. It should be fed until the pigs reach 15lb. body weight. Then they should be fed an approved starter ration.

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite. Do not feed to pigs while they are on any other medicated feed. If signs of toxicity occur, discontinue use of medicated feeds. Not for use in swine weighing more than 250 lb.

Key Information

Protein: 23%

Fat: 7.10%

Lysine: 1.42%

Medication: Denegard/Chlortetracycline

Withdrawal: none

Feed to pigs weighing 8lbs. to 15lbs.

LINDNER 615 PIG STARTER 20% MX BN MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox.....0.0055%(50 grams per ton) Pyrantel Tartrate...0.0106%(96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein-Min.....	20.0%	Lysine-Min.....	1.63%
Crude Fat-Min.....	5.6%	Crude Fiber-Max.....	2.6%
Calcium-Min.....	1.0%	Calcium-Max.....	1.1%
Phosphorous-Min.....	0.64%	Salt-Min.....	None added
Selenium-Min.....	0.4 ppm	Zinc-Min.....	2100 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrus Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Early wean 20% is a complete feed specifically formulated for pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 50lb. body weight. Then they should be fed an approved starter ration.

WARNING: Do not feed to swine weighing more than 75lb. body weight. Do not feed to swine within 10 weeks of slaughter

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

Key Information

Protein: 20%

Fat: 5.6%

Lysine: 1.63%

Medication: MXBN

Withdrawal: 10 weeks

Feed to pigs weighing 15lbs. to 30lbs.

LINDNER 630 PIG STARTER GROWER MX BN MEDICATED

A complete feed for Starting Pigs being fed for Exhibition. For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by Salmonella choleraesuis); aid in the prevention of migration and establishment of large roundworm (Ascaris suum) infections; aid in the prevention of establishment of nodular worm (Oesophagostomum) infections of swine.

ACTIVE DRUG INGREDIENT

Carbadox.....0.0055%(50 grams per ton) Pyrantel Tartrate.....0.0106%(96 grams per ton)

GUARANTEED ANALYSIS

Crude Protein-Min.....	20.0%	Lysine-Min.....	1.3%
Crude Fat-Min.....	4.6%	Crude Fiber-Max.....	2.7%
Calcium-Min.....	0.6%	Calcium-Max.....	0.7%
Phosphorous-Min.....	0.5%	Salt-Min.....	0.2%
Selenium-Min.....	0.4 ppm	Zinc-Min.....	160 ppm

INGREDIENTS

Ground Corn, Ground Oats, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrus Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Pig Starter Grower is a complete feed specifically formulated for pigs fed for exhibition. This product must be fed continuously as the sole ration. It should be fed until the pigs reach 75lb. body weight. Then they should be fed an approved grower finisher ration.

WARNING: Do not feed to swine weighing more than 75lb. body weight. Do not feed to swine within 10 weeks of slaughter

CAUTION STATEMENT

Consult a veterinarian before feeding to severely debilitated animals. Do not use in feeds containing bentonite.

Key Information

Protein: 20%

Fat: 4.6%

Lysine: 1.3%

Medication: MXBN

Withdrawal: 10 weeks

Feed to pigs 30lbs. to 75lbs.

LINDNER 631 STARTER / GROWER TY MEDICATED

A complete feed for Starter/Grower Pigs being fed for exhibition as an aid in the prevention of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery)

ACTIVE DRUG INGREDIENT

Tylosin(50mg/lb)..... 100 grams/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	20.0%	Lysine-Min.....	1.3%
Crude Fat-Min.....	4.6%	Crude Fiber-Max.....	2.7%
Calcium-Min.....	0.6%	Calcium-Max.....	0.7%
Phosphorous-Min.....	0.5%	Salt-Min.....	0.2%
Salt-Max.....	0.3%	Selenium-Min.....	0.4 ppm
Zinc-Min.....	160 ppm		

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrus Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Lindner's Starter Grower TY is a complete feed specifically formulated for pigs fed for exhibition. It should be fed continuously as the sole Medication to pigs weighing not less than 45lb. body weight. Specific feeding instructions may vary according to the goals of the feeding program for show pigs. Generally, pigs can be fed ad libitum or feed may be restricted to achieve desired growth rates.

Key Information

Protein: 20%

Fat: 4.6%

Lysine: 1.3%

Medication: Tylan

Withdrawal: none

Feed to pigs weighing 30lbs. to show weight.

LINDNER

632 STARTER / GROWER LI-FBZ MEDICATED

A complete feed for Starter/Grower Pigs being fed for Exhibition. For reduction in the severity of swine Mycoplasma pneumonia caused by Mycoplasma hyopneumoniae. For the removal of: Lung worms: Metastrongylus apri, Metastrongylus pudendotectus. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (Ascaris suum); nodular (Oesophagostomum dentatum, Ornithostrongylus quadrispinulatum); small stomach (Hyostrongylus rubidus); Adult and larvae (L2, L3, L4 stages intestinal mucosal forms) whipworm (Trichuris suis); Kidneyworm: Adult and larvae Stephanurus dentatus.

ACTIVE DRUG INGREDIENT

Lincomycin (lincomycin hydrochloride) (100mg/lb) 200gm/ton
 Fenbendazole..... 29.2 gm/ton..... (14.6 mg/lb.)

GUARANTEED ANALYSIS

Crude Protein-Min.....20.0%	Lysine-Min.....1.3%	Crude Fat-Min..... 4.6%
Crude Fiber-Max..... 2.7%	Calcium-Min..... 0.6%	Calcium-Max..... 0.7%
Phosphorous-Min..... 0.5%	Salt-Min.....0.2%	Salt-Max.....0.3%
Selenium-Min.....0.3 ppm	Zinc-Min.....160 ppm	

INGREDIENTS: Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrus Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Feed as the sole ration to provide a total intake of 9mg Fenbendazole/kg of body weight (4.08mg Fenbendazole/lb. of body weight) divided over 3-12 days. Multiply 0.28 times pound of body weight and feed this amount of product over 3-12 days. Example:

Body wt of Pig to be Dewormed	Feed Required per pig	Days Req to Consume Dose
50 lb.	14.0 lb.	7
75 lb.	21.0 lb.	7
100 lb.	28.0 lb.	7

CAUTION STATEMENT Occasionally, swine fed Lincomycin my, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. these conditions have been self-correcting within 5-8 days without discontinuing the Lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to seine that weigh more than 250 lbs. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasites.

WARNING: Do not slaughter swine for human consumption for 6 days following last treatment with lincomycin.

Key Information

Protein: 20%

Fat: 4.6%

Lysine: 1.3%

Medication: Lincomix, Safe-Guard

Withdrawal: 6 days

Feed to pigs for 21 days.

LINDNER 633 STARTER / GROWER CSP MEDICATED

A complete feed for Starter / Grower Pigs being fed for Exhibition. Administer to swine in a complete feed for reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by Salmonella choleraesuis and vibronic dysentery); maintenance of weight gains in the presence of atrophic rhinitis; increased rate of weight gain and improved feed efficiency from 10 lb. bodyweight up to 6 weeks postweaning. For swine raised in confinement (dry lot) or on limited pasture.

ACTIVE DRUG INGREDIENT

Chlortetracycline100 gm/ton Sulfathiazole.....100 gm/ton
Penicillin 50 gm/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	20.0%	Lysine-Min.....	1.3%
Crude Fat-Min.....	4.6%	Crude Fiber-Max.....	2.7%
Calcium-Min.....	0.6%	Calcium-Max.....	0.7%
Phosphorous-Min.....	0.5%	Salt-Min.....	0.2%
Salt-Max.....	0.3%	Selenium-Min.....	0.3 ppm
Zinc-Min.....	160 ppm		

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Lindner's Starter Grower ASP is a complete feed specifically formulated for pigs fed for exhibition. It should be fed continuously as the sole medicated ration to pigs weighing up to 75 lb. body weight. Specific feeding instructions may vary according to the goals of the feeding program for show pigs. The animals should be self fed.

CAUTION STATEMENT

WARNING: Withdraw 7 days prior to slaughter.

Key Information

Protein: 20%

Fat: 4.6%

Lysine: 1.3%

Medication: CSP

Withdrawal: 7 days

LINDNER 670 GROWER TY MEDICATED

A complete feed for Grower Pigs being fed for exhibition as an aid in the prevention of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery)

ACTIVE DRUG INGREDIENT

Tylosin (50mg/lb) 100 grams/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	18.0%
Lysine-Min.....	1.10%
Crude Fat-Min.....	4.1%
Crude Fiber-Max.....	2.2%
Calcium-Min.....	0.5%
Calcium-Max.....	0.6%
Phosphorous-Min.....	0.4%
Salt-Min.....	0.2%
Salt-Max.....	0.3%
Selenium-Min.....	0.3 ppm
Zinc-Min.....	160 ppm

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Lindner's Grower TY is a complete feed specifically formulated for pigs fed for exhibition. It should be fed continuously as the sole Medication to pigs weighing 70lb. body weight to show or finish. specific feeding instructions may vary according to the goals of the feeding program for show pigs. Generally, pigs can be fed ad libitum or feed may be restricted to achieve desired growth rates.

Key Information

Protein: 18%

Fat: 4.1%

Lysine: 1%

Medication: Tylan

Withdrawal: none

Feed to pigs 75lbs. to show weight.

LINDNER 671 GROWER BMD-FBZ MEDICATED

A complete feed for Grower Pigs being fed for exhibition. For control of swine dysentery associated with *Brachyspira hyodysenteriae* on premises with a history of swine dysentery but where signs of the disease have not yet occurred or following an approved treatment of the disease condition.

FOR THE REMOVAL OF: Lungworms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular (*Oesophagostomum denatum*, *Ornithostrongylus quadrispinulatum*); small stomach (*Hyostrongylus rubidus*); Adult and larvae (I2, L3, L4 stages-intestinal mucosal forms) whipworm (*Trichuris suis*). Kidney worm: Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENT

Bacitracin Methylene Disalicylate.....250 g/ton Fenbendazole.....29.2 mg/ton(14.6 mg/lb)

GUARANTEED ANALYSIS

Crude Protein-Min.....	18.0%	Lysine-Min.....	1.10%
Crude Fat-Min.....	4.6%	Crude Fiber-Max.....	2.3%
Calcium-Min.....	0.6%	Calcium-Max.....	0.7%
Phosphorous-Min.....	0.5%	Salt-Min.....	0.2%
Salt-Max.....	0.3%	Selenium-Min.....	0.3 ppm
Zinc-Min.....	160 ppm		

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferric Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Feed as a sole ration in a complete feed for a period of 3-12 days to provide a total intake of 9mg fenbendazol/kg of body weight. No withdrawal time needed.

AMOUNT OF NEEDED PER PIG

Feed at the rate of 1lb. complete feed for each 25lb. body weight each day for 7 days.

CAUTION STATEMENT

Not for the use in swine that weigh more than 250lb. Under conditions of continued exposure to parasites, re-treatment may be needed after 4-6 weeks. Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory

Key Information

Protein: 18%

Fat: 4.1%

Lysine: 1%

Medication: Safe-Guard, BMD

Withdrawal: none

Feed to pigs for 3 to 12 days.

LINDNER 674 PIG FINISHER PAYLEANE MEDICATED

A complete feed for Finisher Pigs Being fed for exhibition. For improved feed efficiency and increased carcass leanness, and for the prevention and /or control of porcine proliferative enteropathies (ileitis) associated with Lawsonia intracellularis.

ACTIVE DRUG INGREDIENT

Ractopamine HCl	18.0 g/ton
Tylosin..... (50 mg/lb).....	100 gm /ton

GUARANTEED ANALYSIS

Crude Protein-Min.....18.0%	Lysine-Min.....1.10%
Crude Fat-Min.....4.1%	Crude Fiber-Max.....2.2%
Calcium-Min.....0.5%	Calcium-Max.....0.6%
Phosphorous-Min.....0.4%	Salt-Min.....0.2%
Salt-Max.....0.3%	Selenium-Min.....0.3 ppm
Zinc-Min.....160 ppm	

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysin, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite.

FEEDING DIRECTIONS

Feed continuously to finishing swine as sole ration for 21 days between 150 lb. (68 kg) to 240 lb.(109 kg) bodyweight. Ractopamine HCl should be fed in a complete swine ration containing at least 16% crude protein.

CAUTION: Pigs fed Ractopamine HCl are at an increased risk for exhibiting the downer pig syndrome (also referred to as “slows,” “subs” or “suspects”). Pigs handling methods to reduce the incidence of downer pigs should be thoroughly evaluated prior to initiating use of Ractopamine HCl. Not for use in breeding swine.

Key Information

Protein: 18%

Fat: 4.1%

Lysine: 1%

Medication: Tylan-Ractopamine

Withdrawal: none

Feed 5lbs. of Lindner 674 to get 18gm of intake.

LINDNER 677 GROWER FINISHER LI MEDICATED

A complete feed for Grower/Finisher Pigs being fed for exhibition. For control of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by Lawsonia intracellularis.

ACTIVE DRUG INGREDIENT

Lincomycin (lincomycin hydrochloride)40 gm/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	18.0%	Lysine-Min.....	1.20%
Crude Fat-Min.....	5.0%	Crude Fiber-Max.....	3.2%
Calcium-Min.....	0.7%	Calcium-Max.....	0.75%
Phosphorous-Min.....	0.4%	Salt-Min.....	0.2%
Salt-Max.....	0.3%	Selenium-Min.....	0.4 ppm
Zinc-Min.....	60 ppm		

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferric Sulfate, Defluorinated Phosphate, Mineral Oil, Niacine Supplement, Manganese Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotin, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D 3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Feed as a sole ration. Limitations For Use: For use in animals or on premises with a history of dysentery, but where symptoms have not yet occurred.

CAUTION STATEMENT

Occasionally, swine fed Lincomycin may, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5-8 days without discontinuing the Lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lbs.

Key Information

Protein: 18%

Fat: 5%

Lysine: 1.1%

Medication: Lincomix

Withdrawal: none

Feed to pigs weighing 75lbs. to show weight.

LINDNER 685 SHOW 16% TY MEDICATED

A complete feed for increased rate of weight gain and improved efficiency for grower & finishing pigs.

ACTIVE DRUG INGREDIENT

Tylosin (10mg/lb) 20 grams/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	16.0%
Lysine-Min.....	1.0%
Crude Fat-Min.....	5.0%
Crude Fiber-Max.....	3.0%
Calcium-Min.....	0.6%
Calcium-Max.....	0.7%
Phosphorous-Min.....	0.4%
Salt-Min.....	0.2%
Salt-Max.....	0.3%
Selenium-Min.....	0.3 ppm
Zinc-Min.....	175.0 ppm

FEEDING DIRECTIONS

Lindner's Grower Finisher TY is a complete feed and should be fed continuously to pigs weighing 70 to 220 pounds. Provide fresh water at all times.

CAUTION

Follow feeding directions at all times. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 milligrams per head per day. Store in a dry area.

DO NOT use any product that has become old, molded or insect contaminated.

Key Information

Protein: 16%

Fat: 5.1%

Lysine: 1.08%

Medication: Tylan

Withdrawal: none

Feed to pigs weighing 75lbs. to show weight

LINDNER 686 DIMENSION MEAL MEDICATED

A Complete Feed for Growing Pigs Being Fed for Exhibition
For control of swine dysentery associated with *Brachyspira hyodysenteriae* in pigs up to 250 lbs body weight on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

ACTIVE DRUG INGREDIENT

Bacitratin Methylene Disalicylate..... 250 g/ton

GUARENTEED ANALYSIS

Crude Protein—Min.....14.0%	Lysine—Min..... 0.80%
Crude Fat —Min.....8.2%	Crude Fiber —Max..... 4.3%
Calcium —Min.....0.4%	Calcium —Max.....0.5%
Phosphorus —Min.....0.3%	Salt —Min.....0.3%
Salt —Max.....0.4%	Selenium—Min.....0.3 ppm
Zinc —Min.....162 ppm	

INGREDIENTS

Ground Corn, Soybean Meal, Dried Whey, Animal Fat, Dicalcium Phosphate, Fish Meal, Calcium Carbonate, Salt, L-Lysine, Copper Sulfate, Artificial Flavors, DL-Methionine, Zinc Oxide, Ferrous Sulfate, Defluorinated Phosphate, Mineral Oil, Niacin Supplement, Manganous Oxide, Vitamin E Supplement, Calcium Pantothenate, Magnesium Oxide, Biotine, Menadione Dimethylpyrimidinol Bisulfite, Vitamin A Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B 12 Supplement, Folic Acid, Potassium Iodide, Vitamin D3 Supplement, Sodium Selenite

FEEDING DIRECTIONS

Lindner's Shape & Guts is a complete feed for show pigs from 140 lb to show, and should only be fed to pigs that need more cover and body. Feed as the sole ration. Provide fresh water at all times.

CAUTION

Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 mg. per head per day. Store in a dry area. DO NOT use any product that has become old, molded or insect contaminated.

Key Information

Protein: 14%

Fat: 8.2%

Lysine: .80%

Medication: BMD

Withdrawal: none

Feed to pigs weighing 140lbs. to show weight

LINDNER 690 18% SOW RATION

For Lactating Sows and Gilts

GUARANTEED ANALYSIS

Crude Protein-Min.....	18.10%
Lysine.....	0.81%
Crude Fat-Min.....	3.00%
Crude Fiber-Max.....	3.05%
Calcium-Min.....	0.68%
Calcium-Max.....	0.76%
Phosphorous-Min.....	0.73%
Salt-Min.....	0.33%
Salt-Max.....	0.42%
Selenium-Min.....	0.30 ppm
Zinc-Min.....	204 ppm

FEEDING DIRECTIONS

Feed 4-5 pounds per day during first 2/3 of gestation period. Increase to 6 pounds for last 1/3 of the gestation.

CAUTION

Follow feeding directions at all times. This product contains added selenium. Intake of supplemental selenium should not exceed 0.3 parts per million on a complete diet basis or a total of 3 milligrams per head per day. Store in a dry area.

DO NOT use any product that has become old, molded or insect contaminated.

Key Information

Protein: 18%

Fat: 3.1%

Lysine: 0.9%

Medication: none

Withdrawal: none

Feed 4lbs. per head to breeding gilts and sows

1103

Protein: 20%
 Fat: 5%
 Lysine: 1.5%
 Medication: Ractopamine
 Withdrawal: none
 4oz= 9gm, 8oz= 18gm

Other Feeds

**690
Oat Pellet**

Protein: 12%
 Fat: 4%
 Medication: none
 Withdrawal: none
 Feed to pigs as filler

1104 SUP RRR Oink

Crude Protein.....	23.00 %
Crude Fat.....	30.00%
Crude Fiber.....	0.15%
Calcium / min.....	0.50% / MAX 1.00%
Phosphorous.....	0.65%
Copper.....	.2ppm / MAX 6ppm
Selenium.....	0.3 ppm
Ash.....	6.50%
Lactose, Dry Basis Max.....	25.00%
Moisture Max.....	6.00%
Vit A.....	30,000IU/lb
Vit D.....	10,000IU/lb
Vit E.....	150 IU/lb

1104 is a top dress to be used as a fat supplement to improve the body condition and performance of exhibition swine.

LINDNER 201 SHOW GOAT RU MEDICATED

For Weaned Goats Being Fitted For Show
For the prevention of coccidiosis by Eimeria crandallis,

ACTIVE DRUG INGREDIENT

Monensin.....20 g/ton

GUARANTEED ANALYSIS

Crude Protein-Min.....	17.0%
(This includes not more than 0.8% equivalent crude protein from Non-protein nitrogen)	
Crude Fat-Min.....	3.0%
Crude Fiber-Max.....	11.0%
Calcium-Min.....	0.68%
Calcium-Max.....	0.77%
Phosphorous-Min.....	0.4%
Salt-Min.....	0.04%
Salt-Max.....	0.54%
Copper, Min.....	30ppm
Copper Max.....	40ppm
Selenium-Min.....	0.3000 ppm
Vitamin A, Min.....	10,000 IU per lb.
Vitamin D3, Min.....	5,000 IU per lb.
Vitamin E, Min.....	20 IU per lb
Zinc-Min.....	160 ppm

INGREDIENTS

Ground Corn, Wheat Middlings, Rolled Barley, Dehydrated Alfalfa Meal, Dehulled Soybean Meal, Cane Molasses, Brewers Dried Yeast, Fish Meal, Salt, Calcium Carbonate, Ammonium Chloride Monocalcium Phosphate, Dicalcium Phosphate, natural and Artificial Flavors, Manganous Oxide Zinc Oxide, Vitamin E Supplement, Potassium Sulfate, Magnesium Oxide, Defluorinated Phosphate, mineral Oul, Copper Sulfate, Ferrous Sulfate, Vitamin A Supplement, Vitamin D3 Supplement, Potassium Iodide, Cobalt Carbonate, Sodium Selenite.

FEEDING DIRECTIONS

201 Show Goat RU is a nutritionally balanced ration for wethers and does being fitted for show. Feed complete feed continuously as the sole ration. Note: Always provide clean, fresh water to goats at all times. Goats should be kept parasite free and current on all vaccinations and supplementations recommended by your veterinarian.

CAUTION STATEMENT

Consumption of this product by sheep may result in copper toxicity. Do not allow horses or other equine access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated feed is intended for use in cattle or in goats. Consumption by unapproved species may result in toxic reactions. Do not feed to lactating goats.

Key Information

Protein: 17%

Fat: 3.0%

Fiber: 11%

Medication: Monensin

Withdrawal: none

Free choice salt must be made available to goats

*Committed to our
Customers!*



*Focused on
Results!*

www.lindnerfeed.com

**Lindner Feed & Milling
PO Box 767
Comfort, TX 78013
Mill: 830-995-2143**

**Please direct questions and inquiries to
Conda Maze at 281-642-4017.**

Catalog produced by

